arXiv:astro-ph/0505085v1 4 May 2005

Magnetic White Dwarfs from the SDSS II. The Second and Third Data Releases[1]

Karen M. Vanlandingham[2,3], Gary D. Schmidt[2], Daniel J. Eisenstein[2], Hugh C. Harris[4], Scott F. Anderson[5], Pat B. Hall[6], James Liebert[2], Donald P. Schneider[7], Nicole M. Silvestri[5], Gregory S. Stinson[5], and Michael A. Wolfe[5]

kvanland@as.arizona.edu

ABSTRACT

Fifty-two magnetic white dwarfs have been identified in spectroscopic observations from the Sloan Digital Sky Survey (SDSS) obtained between mid-2002 and the end of 2004, including Data Releases 2 and 3. Though not as numerous nor as diverse as the discoveries from the first Data Release, the collection exhibits polar field strengths ranging from 1.5 MG to ∼1000 MG, and includes two new unusual atomic DQA examples, a molecular DQ, and five stars that show hydrogen in fields above 500 MG. The highest-field example, SDSS J2346+3853, may be the most strongly magnetic white dwarf yet discovered. Analysis of the photometric data indicates that the magnetic sample spans the same temperature range as for nonmagnetic white dwarfs from the SDSS, and support is found for previous claims that magnetic white dwarfs tend to have larger masses than their nonmagnetic counterparts. A glaring exception to this trend is the apparently low-gravity object SDSS J0933+1022, which may have a history involving a close binary companion.

Subject headings: white dwarfs — stars : magnetic fields

---

[1] A portion of the results presented here were obtained with the MMT Observatory, a facility operated jointly by The University of Arizona and the Smithsonian Institution.

[2] Steward Observatory, The University of Arizona, Tucson AZ 85721.

[3] Department of Astronomy, Columbia University, 550 West 120th Street, New York, NY 10027.

[4] U.S. Naval Observatory, P.O. Box 1149, Flagstaff, AZ 86002-1149.

[5] Department of Astronomy, University of Washington, Box 351580, Seattle, WA 98195-1580

[6] Department of Physics & Astronomy, 128 Petrie Science & Engineering Building, York University, 4700 Keele St., Toronto, ON, M3J 1P3, Canada

[7] Pennsylvania State University, Department of Physics & Astronomy, 525 Davey Lab., University Park, PA 16802.

1. Introduction

In October 2004, the release of the third installment from the Sloan Digital Sky Survey (SDSS DR3) brought the database current through June 2003, with images to $g > 22$ for nearly 5300 square degrees of the sky and fiber spectroscopy for selected targets over almost 4200 square degrees (Abazajian et al. 2004; 2005). Though designed principally to probe the extragalactic universe, the survey is providing a wealth of information on stellar populations both on and off the main sequence, as well as an effective discovery tool for stellar objects such as accretion binaries, brown dwarfs, and other classes of unusual stars. In this paper we focus on the magnetic white dwarfs in the SDSS database.

The study of magnetic white dwarfs is important not only because it is possible to detect magnetism in this stage over nearly 6 orders of magnitude in strength, but it is generally thought that the fields on white dwarfs are amplified versions of what pervaded the main sequence progenitors. Magnetic white dwarfs may therefore prove to be critical in understanding the role(s) of magnetism in earlier phases of stellar evolution, and provide a clue to the reason behind the essentially universal presence of magnetic fields on neutron stars. Finally, magnetic white dwarfs provide our only means of empirically studying the effects of fields $B \gtrsim 10$ MG on atomic and molecular emission processes. Indeed, in virtually all cases, observational discovery of magnetically shifted features on white dwarfs has preceded the calculations that explain the splitting. Wickramasinghe & Ferrario (2000) provide an excellent review of the state of field in the pre-SDSS era and amplify on the relevance to broader questions of astrophysics.

The first paper in a series reporting the identification of isolated (non-accreting) magnetic white dwarfs from the SDSS (Schmidt et al. 2003, hereafter Paper I) presented 53 new stars through the first data release (DR1), with magnetic fields in the range $1.5 < B_p < 560$ MG (MG $= 10^6$ G). That list nearly doubled the previously known sample of magnetic white dwarfs and included 3 new magnetic DB (helium feature) stars as well as several with exotic/unknown atmospheric compositions. The current paper reports an additional 52 discoveries from survey data available through the end of 2004, including both DR2 and DR3.

2. Observational Data

The SDSS is compiling an enormous imaging and spectroscopic data set for selected regions of the northern sky using a special-purpose 2.5 m telescope at Apache Pt., New Mexico (e.g., Fukugita et al. 1996; Gunn et al. 1998; Lupton et al. 1999; York et al.

2000; Lupton et al. 2001; Pier et al. 2003). Magnetic white dwarfs are recognized in the spectroscopic database, which targets objects selected from deep, 5-color (u, g, r, i, z) images (Hogg et al. 2001; Smith et al. 2002; Ivezic et al. 2004). Spectroscopic target selection is optimized for QSOs and galaxies, and utilizes complex color criteria that have evolved somewhat over time. This and the highly unusual spectra of strongly magnetic white dwarfs imply that candidate spectra might be found in several target classes. Indeed, based on their photometric colors, the objects reported here were generally selected as QSO (∼40%) or HOT_STD (∼20%), with the remainder sprinkled among STAR_WHITE_DWARF, STAR_BHB, and other categories lying off the stellar locus such as SERENDIPITY_BLUE and SERENDIPITY_DISTANT (see Stoughton et al. 2002 for target category descriptions). The spectroscopy is performed with twin dual-beam spectrographs covering the regions $3900 - 6200$ Å and $5800 - 9200$ Å and providing a resolving power ($\lambda/d\lambda \sim 1800$) sufficient to detect a Zeeman triplet at Hα for $B \gtrsim 1.5$ MG.

Additional optical spectroscopy and spectropolarimetry were obtained during the period Feb.−Dec. 2004 for several unusual and questionable targets. These observations utilized the instrument SPOL (Schmidt et al. 1992) attached to the Steward Observatory 2.3 m Bok telescope on Kitt Peak and the 6.5 m MMT atop Mt. Hopkins. In the configuration used, they provide a spectral coverage of $\sim\lambda\lambda 4200 - 8400$ and resolution ∼15 Å. Details of the observing rationale can be found in Paper I.

3. Spectroscopic Identification

For Paper I, the recognition of magnetic objects among the many thousands of survey spectra was carried out by eye, searching among a variety of target categories of stars and unusual objects. Additional candidate magnetic objects were contributed by workers in other areas of SDSS research. With subjective inspection playing such a central role, questions of completeness arose, questions that were evaluated in part by cross-referencing the SDSS discoveries with previously-known magnetic white dwarfs in surveyed portions of the sky.

Beginning with this edition, the strictly visual identification process was augmented by an automated process originally developed to compile the SDSS DR3 white dwarf catalog (Eisenstein et al. 2005). While that catalog was not specifically designed to find magnetic white dwarfs, it recovered a reasonable number of examples. Briefly, the automated procedure begins with a color cut in $u - g$ and $g - r$ for point sources that have spectra not confidently classified as extragalactic by the SDSS automated software. The spectra and photometry are then fit to a temperature and surface gravity grid of (non-magnetic) pure hydrogen and helium white dwarf atmosphere spectra using the `autofit` program (Kleinman et al. 2004). Deviations from the best-fitting model are computed for a variety of lines, including Ca II λ3933, He I λ4471, He II λ4686, Hβ, Mg Ib λ5184, Hα, and the (1−0) C_2 Swan band near 5165 Å. Stars that present statistically significant deviations at these wavelengths or fail one of a number of other quality assurance tests, such as an acceptable value for the overall reduced χ^2, are flagged for visual inspection. Stars with moderate to strong fields tend to trigger inspections either because of poor overall fits or because of the above line tests. Weak magnetic fields (conservatively, those with $B \lesssim 3$ MG, depending on object brightness) may not perturb the spectra enough for the automated classifier to detect the deviation; they remain among the non-magnetic DA and DB lists and require visual recognition for discovery. We emphasize that in no case does the automated process classify a star as magnetic; rather a spectrum is simply indicated for future visual examination. Furthermore, the automated procedure only contributes to the eventual catalog of magnetic stars; more than half of the discoveries reported here originated from visual assessment of survey spectra, an effort that is continuously under way.

4. Results

4.1. Atmospheric Composition and Field Strength

All but one of the 52 magnetic white dwarfs found in the SDSS database since Paper I show hydrogen features, as listed in Table 1. Fifty are apparently strictly DAs, two are DQAs showing additional lines of atomic carbon, and one shows molecular carbon features probably related to LHS 2229 and SDSS J1333+0016 (Paper I)[1]. The full SDSS coordinate designation (*hhmmss.ss*$\pm$*ddmmss.s*, J2000) is provided in Table 1, though stars are abbreviated by the first four digits of each coordinate in the following text and figures. The table also contains the Plate-MJD-Fiber spectroscopic identifier, International Atomic Time (TAI) date and time of the midpoint of the spectroscopic observation, and comments regarding spectral appearance, data release of the fiber spectrum, followup polarimetry, aliases, etc. Polar field strengths have been determined by visual comparison to model magnetic spectra as described in Paper I. To be consistent, a centered dipole has been assumed for the field structure, even in cases where the fits are obviously not well-represented by this morphology, and the inclination represents the viewing angle relative to the dipole axis, with a typical uncertainty of 30°. Uncertainties in the inclination are larger for very faint or weak-line objects, and inclinations are not quoted at all for fields $\gtrsim$500 MG, where the often very smeared spectra reduce the number of points of detailed comparison.

---

[1] Beyond these, three stars reported in Paper I were rediscovered from new targetings of previously-observed SDSS fields. In Plate-MJD-Fiber designation, these are 710-52203-311 = 709-52205-120 = 411-51817-172 = SDSS J0304−0025, 2049-53349-450 = 810-52326-392 = 415-51810-370 = SDSS J0331+0045 = KUV 03292+0035, and the magnetic DB star 1898-53260-299 = 429-51820-311 = SDSS J0142+1315. Two stars contained in Table 1 are rediscoveries of white dwarfs not previously recognized as magnetic.

Derived polar field strengths for the best-fitting dipole range from 1.5 MG to ~1000 MG, and quoted values can generally be considered accurate to better than 10%. The current list is distinguished by no less than 5 examples with hydrogen features at $B_p > 500$ MG: SDSS J0021+1502 (550 MG); SDSS J1351+5419 (760 MG); SDSS J1206+0813 (~830 MG); SDSS J1003+0538 (900 MG), and SDSS J2346+3853 (1000 MG). The latter star rivals PG 1031+234 (Schmidt et al. 1986) for the highest field strength yet detected on a white dwarf. SDSS J1351+5419 is a rediscovery of a previously-known magnetic star from the Second Byurakan objective-prism QSO survey (Liebert et al. 1994). Spectra of the new high-field DAs comprise an instructive progression with field strength, as shown in Figure 1. For $B_p \gtrsim 400$ MG, persistent lines appear at ~4200 Å and 4550 Å, formed at near-stationary points of the Hβ transitions $2s0$ to $4f0$ and $2s0$ to $4f{-1}$, respectively. Other distinctive features at high fields are the pair of Hα components ($2s0$ to $3p0$ and $2p{-1}$ to $3d{-2}$) that slowly cross in the region ~6000 − 7000 Å, and the stationary $2p0$ to $3d{-1}$ component of Hα at ~8500 Å can be recognized in the brighter stars.

The existence of strong magnetic fields on SDSS J2346+3853, SDSS J0021+1502, and SDSS J1206+0813 was confirmed through circular (and in the former case linear) spectropolarimetry. At such strong fields, polarization is more a property of the continuum than the lines (e.g., Schmidt et al. 1996), so the polarization spectra are not highly structured, and the values entered in Table 1 are averages over the range $\lambda\lambda 4200 - 8400$. The modest polarization of SDSS J2346+3853, when compared with other very strongly magnetic DA white dwarfs like PG 1031+234 (Schmidt et al. 1986), Grw +70°8247 (Angel et al. 1985), or SDSS J2247+1456 (Paper I), suggests that the field morphology on the new star may be unusually tangled.

Spectra of the two magnetic DQA stars are presented in Figure 2. SDSS J0236−0808 was observed as part of DR1 but was not recognized as magnetic in Paper I. The object is faint, the line splitting is only partially resolved at $B_p = 5$ MG, and the presence of carbon features probably confused the visual identification. The rather weak magnetic field is consistent with an overall lack of circular polarization ($v = V/I < 0.1\%$). SDSS J1328+5908 was included among the 18 carbon-feature white dwarfs discovered in the SDSS through 2003 (Liebert et al. 2003b). Here we point out that the hydrogen lines appear as clear Zeeman triplets in a modest magnetic field, as indicated in the figure for $\langle B \rangle = 8.5$ MG. The spectra of C I and C II have not yet been calculated in the Paschen-Back regime, but at this field strength the transitions can be assumed to resemble normal triplets, with any given ensemble possibly somewhat skewed and displaced to the blue by the quadratic Zeeman effect. The π components of these features would be expected to line up near zero-field wavelengths, and indeed there is a good correspondence between several of the observed features and the principal lines observed in atomic DQ white dwarfs (marked in Fig. 2, as taken from Liebert et al. 2003b).

SDSS J0954+0913 is an interesting example of a white dwarf dominated by molecular absorption that may represent a "transition" object between normal (C_2) DQ stars and the "peculiar" DQs discussed by Schmidt et al. (1995). A comparison is made of an MMT spectrum of SDSS J0954+0913 obtained with SPOL on 2004 24 April 24 with spectra of SDSS J2310−0057 (DQ; Harris et al. 2003) and LHS 2229 (magnetic peculiar DQ; Schmidt et al. 1999) in Figure 3. Both the positions and shapes of the bands in SDSS J0954+0913 are intermediate between the other two stars, suggesting that the features in all three are the C_2 Swan bands. If so, a C_2 identification (Bues 1999) for the displaced and distorted bands in LP 790-29 (and SDSS J1113+0146) might be called into question, since these lack the "scalloped" appearance of the former stars. The spectrum-added circular polarization for SDSS J0954+0913, obtained from the same MMT spectropolarimetric data, is $v = -0.93\%$. This is significant but less in magnitude than the −3.8% found for LHS 2229. The polarization within the bands approaches +20% in the latter object, but no such increase is seen in SDSS J0954+0913. If, as has been suggested, LHS 2229 has a surface magnetic field of ∼100 MG, it would appear that the value on SDSS J0954+0913 is substantially less.

Three additional objects whose survey spectra are displayed in Figure 4 deserve mention here. SDSS J1007+1237 exhibits clear Zeeman patterns in its Balmer series, but the structure in at least Hα and Hβ is more complex than the model spectra predict for a polar field strength of ∼7 MG. The spectral energy distribution is also very odd, suggesting either a calibration mismatch between the blue and red spectrograph channels or a broad depression extending from ∼5800 Å to at least Hα. Spectropolarimetric observations are planned of this faint target to investigate whether it might be a magnetic white dwarf pair or a single star with very complex field structure. The second star, SDSS J1234+1248, also shows Zeeman triplets in a field of ∼7 MG, but the π components are both unusually sharp and considerably deeper than the models predict; indeed deeper than any other low-field DA in the sample. Possible interpretations include a magnetic/nonmagnetic DA pair (DAH+DA), or once again a distinctly nondipolar field geometry. In this case a large portion of the star would be covered by very weak magnetic field ($\lesssim$1 MG). An example of analogous field structure at a somewhat weaker strength is WD 1953−011 (Maxted et al. 2000). Finally, SDSS J9044+5321 presents a challenge. The Balmer lines are seen at their rest wavelengths, however other features are clearly visible in the spectrum. This is strong evidence of a binary system. Attempt to match the additional lines with DQ or DB features have been unsuccessful. A magnetic DA model with a polar field near 53 MG will cause the π component of Hα to shift enough to match up with the blue wing on the observed profile but the model is unable to reproduce the features seen in the Hβ region. Followup observations are needed to determine the true nature of this system.

4.2. Photometry and Stellar Parameters

In Table 2 we collect the *psfmag* photometry of the new magnetic stars from survey imaging data. The final column contains a temperature estimate derived from a comparison of a star's observed colors (primarily $u-g$ and $g-r$) to those computed from spectral models for nonmagnetic DA white dwarfs with various values of $\log g$ by Bergeron et al. (1995). No temperature is listed for SDSS J0954+0913 because the optical spectral energy distribution is strongly affected by the deep molecular bands. For SDSS J1508+3945, the photometry led to inconsistent temperature estimates among the colors; indeed the *psfmag* values do not track the fiber spectrum and differ by up to 0.6 mag from the "fiber" magnitudes. In this case the quoted temperature is based on the fiber magnitudes, which are provided in the table footnote. Temperatures derived from broad-band colors are prone to errors because of a magnetic field's effect on the continuum opacity and absorption lines, both of which increase in importance with field strength. Values whose uncertainties are estimated to be larger than 1000 K, either because of a strong field or because they lie off the locus in the model 2-color diagram, are flagged by a colon in the table. Derived temperatures span 7,000 K to ∼28,000 K, thus occupying the same range as the bulk of nonmagnetic SDSS DA white dwarfs from Kleinman et al. (2004), whose temperatures and gravities were derived by line-profile fitting. Unfortunately, the cool white dwarfs that comprise the peak of the white dwarf luminosity function are not targeted by the survey unless they have unusual spectral energy distributions. This selection effect likely contributes to the fact that nearly all of the magnetic discoveries from the SDSS are of DA spectral type.

While the 2-color diagrams are most useful for temperature estimation, the $u-g, g-r$ plot also offers some leverage on the stellar surface gravity through the sensitivity of the $u-g$ color to the Balmer discontinuity (see, e.g., Figure 1 of Harris et al. 2003). Estimates for $\log g$ based on broadband colors are not as accurate as those from the popular method of fitting Balmer-series line profiles (Bergeron et al. 1992), but the spectroscopic technique is rendered ineffective by the reduction in Stark broadening in strong magnetic fields. Even photometric estimates of $\log g$ for the SDSS magnetic examples are interesting as an ensemble because of mounting evidence that the typical magnetic white dwarf has a higher gravity, and is therefore more massive, than its nonmagnetic counterpart (Liebert et al. 2003a and references therein). The 52 stars from Table 2 that can be used for this purpose bear out this effect, with the average gravity, $\langle \log g \rangle = 8.31$, being higher than the value of 8.06 that resulted from an application of the spectroscopic method to the 1888 nonmagnetic DA white dwarfs from DR1 (Kleinman et al. 2004). The comparison is complicated by the facts that the spectroscopic technique seems to show a bias for high gravities below $T_{\rm eff} = 10,000$ K (Kleinmann et al. 2004), and the photometric method is subject to interstellar reddening. Within the SDSS database, the DR2 and DR3 data are uncorrected for galactic extinction and we have not applied any correction of our own in the course of our analysis. By comparison, the colors used in Kleinman et al. (2004) had full Galactic reddening corrections applied. The differences are generally small, with a median $E(B-V) = 0.034$ mag (http://www.sdss.org/dr3/algorithms/spectrophotometry.html), but we note that applying reddening corrections to the DR2/DR3 magnetic white dwarf photometry would tend to *increase* the inferred gravities for stars hotter than 10,000 K, and we would expect to see an even larger difference between the average gravity of the magnetic and nonmagnetic samples. Either of the above mean gravities are larger than the $\langle \log g \rangle = 7.91$ found by Bergeron et al. (1992) for 129 hot DAs from the McCook & Sion (1987) catalog. The difference corresponds to a typical mass excess for the magnetic stars of $\sim$0.30 $M_\odot$, bringing the mean mass for the SDSS magnetic white dwarfs near the $\sim$0.93 $M_\odot$ derived for the magnetic samples of Liebert et al. (2003a).

In light of the above result, SDSS J0933+1022 is a particularly glaring outlier with a gravity estimate of only $\log g \sim 6.3$. The estimate results from an unusually large $u-g$ color of +0.57, but a low gravity is supported by the lack of detectable wings to the Balmer lines in the survey spectrum shown in Figure 4. Unfortunately, magnetic effects on the profiles invalidates a determination of the gravity by the spectroscopic method, and in any case that technique exhibits an unexplained, systematic problem with fitting DA stars cooler than about 11,000K, as discussed in Kleinman et al. (2004). The 1.5 MG polar field strength of SDSS J0933+1022 is weak for a white dwarf, but sufficient to rule out a main-sequence or horizontal-branch nature, while the temperature of 8,500 K eliminates a subdwarf possibility and at the same time accents the lack of an obvious spectral companion. The star shares some similarities with SDSS J1234−0228, which was modeled by Liebert et al. (2004) to have $\log g = 6.38 \pm 0.05$ and $T_{\rm eff} = 17,500$ K, and interpreted as a helium-composition white dwarf with $M = 0.18 - 0.19$ $M_\odot$ resulting from some type of evolution with a yet undiscovered binary companion. However, considering both its surface temperature and gravity, SDSS J0933+1022 perhaps best resembles the companion to the pulsar J1012+5307, estimated by Van Kerkwijk et al. (1996) and Callanan et al. (1998) to have $T_{\rm eff} \sim 8,600$ K and $M = 0.16$ $M_\odot$. The star demands followup study, both to confirm its low gravity and to look for a massive cool white dwarf, very low-mass main-sequence, or dead pulsar companion.

4.3. Issues of Completeness

Including the additions from this paper, a total of 40 magnetic white dwarfs were identified in the first 1360 $\deg^2$ of the SDSS released as DR1. This should be compared with only 49 new magnetic stars discovered in the additional 2828 $\deg^2$ covered by DR2 and DR3 (22 discoveries were reported in Paper I). There is actually good reason to believe that our current identification procedures are more effective than the strictly visual method of Paper I, once a spectrum is obtained. The machine selection algorithm is not only far more efficient but also more uniform, and the 3 DR1 stars overlooked in the course of preparing Paper I but recovered here attest to its utility. Of course, the areal density of stars varies with location on the sky, but the considerable discrepancy in discovery rate of magnetic stars suggests that improvements in the photometric reduction pipeline and continued refinement of the QSO targeting criteria have reduced the chance that an interloper is selected for a spectroscopic fiber. Comparison of the locations of magnetic white dwarfs in the SDSS color planes for the DR1 and DR2/DR3 samples reveal no regions populated solely by DR1 objects, but the photometric scatter around the locus is found to be slightly reduced in the later sample for the shorter wavelength filter bands. More direct evidence for a reduction in the number of stellar targetings because of improvements in the QSO targeting criteria is provided by the fact that, of the 8 previously-known magnetic white dwarfs that lie in regions of the sky covered by DR2/DR3, only two - SDSS J1214−0234 = LHS 2534 (reported in Paper I) and SDSS J1351+5419 = SBS1349+5434 - were recovered in the survey. Admittedly, G99-37, HS1412+6115, and GD 185 exceed the brightness limit for a spectroscopic fiber, but the three magnetic DAs LHS 2273 (18 MG), PG 1312+098 (10 MG), and G62-46 (7.4 MG) are all within the limits of the SDSS. None was targeted for spectroscopy. This result should be contrasted with the fact that 5 of the 6 qualifying stars in DR1 were recovered and reported in Paper I.

The implied incompleteness of recent (and forthcoming) editions of the survey raises concerns over the effects that a differential selection bias might have on the derived distributions of magnetic stars as functions of field strength or temperature, and thus on inferences that may be drawn from these distributions. However, the breakdown of new identifications from this paper, per decade of field strength, is: 1−10 MG, 13; 10−100 MG, 11; 100−1000 MG, 3. Within the sampling statistics, these are consistent with the corresponding numbers 23:28:9 for stars from Paper I, and with the distribution shown in Figure 8 of that paper for all 116 magnetic white dwarfs known at the time. The question of completeness for magnetic white dwarfs discovered from the SDSS will be revisited when the survey is concluded.

Funding for the creation and distribution of the SDSS Archive has been provided by the Alfred P. Sloan Foundation, the Participating Institutions, the National Aeronautics and Space Administration, the National Science Foundation, the U.S. Department of Energy, the Japanese Monbukagakusho, and the Max Planck Society. The SDSS Web site is http://www.sdss.org/. The SDSS is managed by the Astrophysical Research Consortium (ARC) for the Participating Institutions. The Participating Institutions are The University of Chicago, Fermilab, the Institute for Advanced Study, the Japan Participation Group, The Johns Hopkins University, the Korean Scientist Group, Los Alamos National Laboratory, the Max-Planck-Institute for Astronomy (MPIA), the Max-Planck-Institute for Astrophysics (MPA), New Mexico State University, University of Pittsburgh, University of Portsmouth, Princeton University, the United States Naval Observatory, and the University of Washington. Support for the study of magnetic stars and stellar systems at Steward Observatory is provided by NSF grant AST 03-06080.

REFERENCES

Abazajian, K., et al. 2004, AJ, 128, 502

________. 2005, AJ, in press

Angel, J.R.P., Liebert, J., & Stockman, H.S. 1985, ApJ, 292, 260

Bergeron, P., Saffer, R.A., & Liebert, J. 1992, ApJ, 394, 228

Bergeron, P., Wesemael, F., & Beauchamp, A. 1995, PASP, 107, 1047

Bues, I. 1999, in ASP Conf. Ser. 169, Eleventh European Workshop on White Dwarfs, ed. J.-E. Solheim & E.G. Meistas (San Francisco: ASP), 240

Callanan, P.J., Garnavich, P.M., & Koester, D. 1998, MNRAS, 298, 207

Eisenstein, D.J., et al. 2005, in preparation

Fukugita, M., Ichikawa, T., Gunn, J.E., Doi, M., Shimasaku, K., & Schneider, D.P. 1996, AJ, 111, 1748

Gunn, J.E., et al. 1998, AJ, 116, 3040

Harris, H.C. et al. 2003, AJ, 126, 1023

Hogg, D.W., Finkbeiner, D.P., Schlegel, D.J., & Gunn, J.E. 2001, AJ, 122, 2129

Ivezic, Z., et al. 2004, AN, 325, 583

Kleinman, S.J., et al. 2004, ApJ, 607, 426

Liebert, J., Bergeron, P., Eisenstein, D., Harris, H.C., Kleinman, S.J., Nitta, A., & Krzesinksi, J. 2004, ApJ, 606, L147

Liebert, J., Bergeron, P., & Holberg, J.B. 2003a, AJ, 125, 348

Liebert, J., et al. 2003b, AJ, 126, 252

Liebert, J., Schmidt, G.D., Lesser, M., Stepanian, J.A., Lipovetsky, V.A., Chaffee, F.H., Foltz, C.B., & Bergeron, P. 1994, ApJ, 421, 733

Lupton, R.H., Gunn, J.E., Szalay, A.S. 1999, AJ, 118, 1406

Lupton, R.H., Gunn, J.E., Ivezic, Z., Knapp, G.R., Kent, S.M., & Yasuda, N. 2001, ASP Conf. Ser. 238, p. 269

Maxted, P.F.L., Ferrario, L., Marsh, T.R., & Wickramasinghe, D.T. 2000, MNRAS, 315, L41

McCook, G.P., & Sion, E.M. 1987, ApJ, 244, 280

Pier, J.R., Munn, J.A., Hindsley, R.B., Hennessy, G.S., Kent, S.M., Lupton, R.H., & Ivezic, Z. 2003, AJ, 125, 1559

Schmidt, G.D., Allen, R.G., Smith, P.S., & Liebert, J. 1996, ApJ, 463, 325

Schmidt, G.D., Bergeron, P., & Fegley, B. Jr. 1995, ApJ, 443, 274

Schmidt, G.D., et al. 2003, ApJ, 595, 1101 (Paper I)

Schmidt, G.D., Liebert, J., Harris, H.C., Dahn, C.C., & Leggett, S.K. 1999, ApJ, 512, 916

Schmidt, G.D., Stockman, H.S., & Smith, P.S. 1992, ApJ, 398, L57

Schmidt, G.D., West, S.C., Liebert, J., Green, R.F., & Stockman, H.S. 1986, ApJ, 309, 218

Smith, J.A., et al 2002, AJ, 123, 2121

Stoughton, C. et al. 2002, AJ, 123, 485

Van Kerkwijk, M.H., Bergeron, P., & Kulkarni, S.R. 1996, ApJ, 467, L89

Wickramasinghe, D.T., & Ferrario, L. 2000, PASP, 112, 873

York, D.G. et al. 2000, AJ, 120, 1579

This preprint was prepared with the AAS LaTeX macros v5.2.

Table 1. NEW SDSS MAGNETIC WHITE DWARFS

Star (SDSS+)	Plate-MJD-Fiber	TAI Date	TAI	B_p (MG)	i (deg)	Comments
J002129.00+150223.7	753-52233-432	2001 Nov 20	4:27	550	...	$v = +0.91\%$
J021148.22+211548.2	2046-53327-048	2004 Nov 11	6:13	210	90	post-DR3
J023609.40−080823.9	455-51909-474	2000 Dec 31	3:28	5	90	DQA; $v = +0.01\%$; in DR1
J074850.48+301944.8	889-52663-507	2003 Jan 24	4:54	10	60	
J080440.35+182731.0	2081-53357-442	2004 Dec 18	7:19	49	30	post-DR3
J080502.29+215320.5	1584-52943-132	2003 Oct 30	11:17	5	60	post-DR3
J080938.10+373053.8	758-52253-044	2001 Dec 7	7:55	40	30	
J081648.71+041223.5	1184-52641-329	2003 Jan 2	8:23	10:	30:	
J082835.82+293448.7	1207-52672-635	2003 Feb 2	4:30	30	90:	
J084008.50+271242.7	1587-52964-059	2003 Nov 21	9:35	10	60	post-DR3
J090632.66+080716.0	1300-52973-148	2003 Nov 30	11:31	10	90	post-DR3
J090746.84+353821.5	1212-52703-187	2003 Mar 5	4:28	15	60	
J091124.68+420255.9	1200-52668-538	2003 Jan 23	10:05	45	60	
J091437.40+054453.3	1193-52652-481	2003 Jan 11	9:19	9.5	90	
J093356.40+102215.7	1303-53050-525	2004 Feb 12	9:17	1.5	60:	= SDSS 1303-53047-525; post-DR3
J093447.90+503312.2	901-52641-373	2003 Jan 2	9:38	9.5	60	
J094458.92+453901.2	1202-52672-577	2003 Feb 2	8:24	14	90	
J095442.91+091354.4	1306-52996-004	2004 Dec 23	10:54	...	...	mol. DQ; $v = -0.93\%$; post-DR3
J100356.32+053825.6	996-52641-295	2003 Jan 2	10:50	900:	...	
J100715.55+123709.5	1745-53061-313	2004 Feb 26	4:05	7	60	post-DR3
J101529.62+090703.8	1237-52762-533	2003 May 3	4:50	5:	90:	
J111812.67+095241.4	1222-52763-477	2003 May 4	4:37	6	60:	
J112924.74+493931.9	966-52642-474	2003 Jan 3	10:12	5	60	
J113756.50+574022.4	1311-52765-421	2003 May 6	5:32	9	60:	
J114829.00+482731.2	1446-53080-324	2004 Mar 16	6:00	33	90	post-DR3
J120150.10+614257.0	778-52337-264	2002 Mar 4	8:22	20	90	
J120609.80+081323.7	1623-53089-573	2004 Mar 25	6:24	830:	...	$v = +0.41\%$; post-DR3
J120728.96+440731.6	1369-53089-048	2004 Mar 25	7:53	2.5	90	post-DR3
J122249.14+481133.1	1451-53117-582	2004 Apr 22	5:56	8	90:	post-DR3
J122401.48+415551.9	1452-53112-181	2004 Apr 17	6:30	23:	60	post-DR3
J123414.11+124829.6	1616-53169-423	2004 Jun 13	3:57	7	60:	WD1231+130=LBQ 1231+1305; post-DR3
J124806.38+410427.2	1456-53115-190	2004 Apr 20	4:54	8	90	post-DR3
J125044.42+154957.4	1770-53171-530	2004 Jun 15	4:16	20	60	post-DR3
J125416.01+561204.7	1318-52781-299	2003 May 22	5:54	52	30	

Table 1—Continued

Star (SDSS+)	Plate-MJD-Fiber	TAI Date	TAI	B_p (MG)	i (deg)	Comments
J132002.48+131901.6	1773-53090-011	2004 Mar 26	10:23	5	60	post-DR3
J132858.20+590851.0	959-52411-504	2002 May 17	5:56	18	90	DQA
J135141.13+541947.4	1323-52797-293	2003 Jun 2	7:25	760	20	SBS1349+5434
J142703.40+372110.5	1381-53089-182	2004 Mar 25	11:12	30	60:	post-DR3
J143218.26+430126.7	1396-53112-338	2004 Apr 17	10:17	1.5	90	post-DR3
J143235.46+454852.5	1288-52731-449	2003 Apr 2	7:01	10	30	
J145415.01+432149.5	1290-52734-469	2003 Apr 5	10:26	5	...	
J150813.20+394504.9	1398-53146-633	2004 May 21	8:01	20	90	WD1506+399 = CBS 229; post-DR3
J151130.20+422023.0	1291-52735-612	2003 Apr 6	10:12	12	60	
J164357.02+240201.3	1414-53121-191	2004 Apr 23	11:16	4	90	post-DR3
J164703.24+370910.3	818-52395-026	2002 May 1	9:38	2:	90:	
J165029.91+341125.5	1175-52791-482	2003 Jun 1	9:53	3:	0:	
J170400.01+321328.7	976-52413-319	2002 May 17	10:55	5	90:	
J171556.29+600643.9	354-51792-318	2000 Sep 5	4:27	4.5	60	in DR1
J214900.87+004842.8	1107-52968-374	2003 Nov 22	1:50	10	60	post-DR3
J231951.73+010909.3	382-51816-565	2000 Sep 29	4:44	1.5:	90:	in DR1
J234605.44+385337.7	1883-53265-272	2004 Sep 17	9:22	1000	...	$v = -1.51\%$; $P = 1.33\%, \theta = 164°$; post-DR3
J234623.69−102357.0	648-52559-142	2002 Oct 12	3:56	2.5	90:	

Table 2. PHOTOMETRY OF SDSS MAGNETIC WHITE DWARFS

Star (SDSS+)	u	g	r	i	z	T_{eff} (K)
J002129.00+150223.7	18.69	18.18	17.97	17.93	17.99	7000
J021148.22+211548.2	17.26	17.07	17.21	17.41	17.63	12000
J023609.40−080823.9	19.96	19.76	19.78	19.84	19.95	10000
J074850.48+301944.8	17.39	17.58	17.87	18.49	18.56	22000:
J080440.35+182731.0	18.45	18.13	18.25	18.32	18.55	11000
J080502.29+215320.5	18.25	18.60	18.98	19.37	19.73	28000:
J080938.10+373053.8	19.32	19.02	19.26	19.49	19.76	14000
J081648.71+041223.5	20.74	20.40	20.52	20.63	21.11	11500
J082835.82+293448.7	19.67	19.74	20.05	20.32	20.31	19500
J084008.50+271242.7	19.53	19.18	19.35	19.54	19.87	12250
J090632.66+080716.0	18.62	18.65	18.82	19.11	19.32	17000:
J090746.84+353821.5	19.76	19.63	19.92	20.27	20.48	16500
J091124.68+420255.9	19.17	18.85	18.91	19.02	19.24	10250
J091437.40+054453.3	17.44	17.33	17.64	17.89	18.16	17000
J093356.40+102215.7	19.34	18.77	18.73	18.71	18.73	8500
J093447.90+503312.2	19.19	18.81	18.80	18.85	19.07	8900
J094458.92+453901.2	20.01	19.95	20.14	20.35	20.74	15500:
J095442.91+091352.4	20.07	20.16	19.71	19.67	19.97	...
J100356.32+053825.6	17.96	18.12	18.48	18.65	18.90	23000
J100715.55+123709.5	18.78	18.77	19.05	19.32	19.57	18000
J101529.62+090703.8	19.10	18.59	18.41	18.39	18.49	7200
J111812.67+095241.4	19.17	18.75	18.85	19.04	19.19	10500
J112924.74+493931.9	18.42	18.00	18.08	18.15	18.34	10000
J113756.50+574022.4	17.32	16.85	16.75	16.75	16.77	7800
J114829.00+482731.2	17.88	18.16	18.58	18.84	19.18	27500
J120150.10+614257.0	18.80	18.48	18.57	18.60	18.78	10500
J120609.80+081323.7	19.27	19.04	19.22	19.42	19.58	13000
J120728.96+440731.6	19.44	19.33	19.62	19.97	20.52	16750
J122249.14+481133.1	19.11	18.72	18.72	18.79	18.97	9000
J122401.48+415551.9	19.27	18.96	18.98	19.06	19.25	9500

Table 2—Continued

Star (SDSS+)	u	g	r	i	z	T_{eff} (K)
J123414.11+124829.6	17.85	17.38	17.32	17.36	17.49	8200
J124806.38+410427.2	18.40	17.93	17.71	17.72	17.79	7000
J125044.42+154957.4	18.64	18.26	18.32	18.38	18.20	10000
J125416.01+561204.7	19.36	18.98	19.20	19.33	19.59	13250
J132002.48+131901.6	19.84	19.49	19.82	20.07	20.19	14750
J132858.20+590851.0	17.63	17.85	18.25	18.55	18.89	25000
J135141.13+541947.4	16.66	16.40	16.55	16.72	16.94	12000
J142703.40+372110.5	17.55	17.57	17.91	18.18	18.43	19000
J143218.26+430126.7	18.86	18.99	19.41	19.71	20.15	24000
J143235.46+454852.5	20.06	19.94	20.24	20.49	20.52	16750
J145415.01+432149.5	19.48	19.06	19.21	19.34	19.72	11500
J150813.20+394504.9	17.35	17.90	17.75	18.03	18.51	17000[a]
J151130.20+422023.0	18.20	17.98	18.01	18.19	18.46	9750
J164357.02+240201.3	19.43	19.27	19.58	19.82	20.40	16500
J164703.24+370910.3	17.80	17.64	17.92	18.21	18.50	16250
J165029.91+341125.5	19.13	18.78	18.83	18.98	19.09	9750
J170400.01+321328.7	20.30	20.43	20.83	21.07	21.91	23000
J171556.29+600643.9	19.70	19.35	19.59	19.73	19.78	13500
J214900.87+004842.8	20.12	19.87	19.96	20.19	20.05	11000
J231951.73+010909.3	18.82	18.50	18.44	18.52	18.61	8300
J234605.44+385337.7	18.64	18.89	19.28	19.48	19.85	26000
J234623.69−102357.0	18.77	18.45	18.40	18.44	18.59	8500

[a] Temperature based on *ugriz* “fiber” magnitudes of 17.40, 17.30, 17.60, 17.88, 18.21, respectively.



Fig. 1.— Progression of new strongly-magnetic DA white dwarfs showing features of hydrogen in fields up to 1000 MG, and compared with computed wavelengths for transitions that become stationary or execute turnarounds in the field strength range of interest. Persistent at the highest fields is the pair of Hβ components that become nearly stationary in the range 4200−4600 Å and the two Hα components that wander between 6000−7000 Å. Quoted field strengths are estimated polar values for the best-fit assumed dipole model. Spectra have been smoothed to a resolution of ∼8 Å, scaled, and shifted for display purposes.



Fig. 2.— Two unusual magnetic DQA (hydrogen + carbon) white dwarfs. For SDSS J0236−0808, Zeeman triplets in the Balmer lines are legible by comparison with the theoretical curves shown, with field strength indicated along the right ordinate. The largely blended molecular and atomic carbon features render a somewhat choppy appearance to the spectrum. SDSS J1328+5908 is at a field strength where all features should resemble Zeeman triplets, but carbon has not been calculated in this regime, so we simply point out that the π components of split features should be found near the zero-field wavelengths of the same lines (indicated by tick marks below the observed spectrum). The inset displays the Hγ region for SDSS J0236−0808 at a larger scale, and a horizontal dashed line indicates the zero-flux level for SDSS J1328+5908.



Fig. 3.— Spectral comparison between SDSS J0954+0913, the normal (C_2 Swan band) molecular white dwarf SDSS J2310−0057, and the peculiar magnetic DQ star LHS 2229. At $v = -0.93\%$, circular polarization of SDSS J0954+0913 has been measured to be significant, but it is smaller in magnitude than the $v = 4 - 20\%$ seen in LHS 2229, supporting an interpretation for SDSS H0954+0913 as a intermediate object with a modest magnetic field ($B < 100$ MG). Horizontal dashed lines indicate the zero-flux levels for successive spectra.



Fig. 4.— Survey spectra of four unusual examples. SDSS J1007+1237 and SDSS J1234+1248 both have polar fields $B_p \sim$7 MG, but the former shows a distorted spectral energy distribution, while the latter exhibits conspicuously deep π components suggestive of a nonmagnetic companion or a strongly nondipolar field structure over the surface of the star. We note that the distortion in the spectrum of SDSS J1007+1237 may be due to a calibration mismatch between the blue and red spectrograph channels. The odd Hα profile of SDSS J0944+5321 is suggestive of a companion but DQ, DB, and magnetic DA models have all been unsuccessful in reproducing the features seen in the Hβ region. Photometry of SDSS J0933+1022 implies a very low gravity, $\log g < 7$, suggesting a history involving a close binary companion. Insets display the Hβ and Hα regions for SDSS J0933+1022 at a larger scale, and horizontal dashed lines indicate the zero-flux levels for successive spectra.